

# IDENTIFICATION OF A FAMILY OF ALTERNATIVELY-SPICED ISOFORMS OF ANGIOPOIETIN-1 IN TUMOR CELLS, UPREGULATED BY THROMBIN. Y.-Q. Huang\*, J.-J. Li\*, S. Karpatkin. NYU Medical School, NY, USA.

Angiogenesis is essential for the growth and metastasis of tumors; and thrombin enhances experimental pulmonary metastasis (Blood 90:400a, 1997). We therefore looked for the presence of angiotensin-1 (Ang-1) in tumor cells, a recently-described ligand for the endothelial specific receptor tyrosine kinase, Tie-2 required for embryonic vascular development (Cell 87:1171, 1996); and its possible regulation by thrombin. Ang-1 was detected by RT-PCR and Northern analysis in 4 of 6 cell lines studied: megakaryocyte CHRF-288, breast cancer MDA-MB468, MDA-MB231 and normal fibroblast (FS4); not in breast MCF7 or prostate DU145 cell lines. Thrombin upregulated Ang-1 mRNA in all 4 lines in which it was detected. Ang-2, an inhibitor of Ang-1, was not detected nor induced by thrombin. Surprisingly, three alternatively-spliced species of Ang-1 mRNA (Ang-1.3 Kb, 0.9 Kb and 0.7 Kb of the coding region, in addition to the full length 1.5 Kb) were identified by RT-PCR, cloned and sequenced (fifty clones examined, 8 sequenced). This was confirmed by RT-PCR using specific sense primers derived from junction sites and anti-sense primers from the 3' end of the cDNA for Ang-1. The distribution and/or expression of all 4 isoforms varied in different cells. CHRF had all 4 isoforms confirmed by Northern analysis. MD468 and FS4 had two (1.5 Kb and 1.3 Kb), MD231 had one (1.5 Kb). All 4 isoforms were upregulated and/or induced by thrombin, 0.5-1 u/ml (5-10 nM). Thrombin induction was noted as early as 30 min in FS4 cells and peaked at 3 hrs. Ang-1 contains a secretory peptide, a coiled-coil region thought to be involved in protein assembly into multimeric structures (homo or heteromultimers of related proteins) and a fibrinogen-like region. All four isoforms noted could be synthesized with a hydrophobic leader peptide for secretion, indicating their possible physiologic significance. The coiled-coil region is intact in the 1.5 Kb and 1.3 Kb isoforms, partially deleted in the 0.7 Kb isoform and mostly deleted in the 0.9 Kb isoform. We suggest that the various isoforms described in tumor cells may have different pro- or anti-angiogenic functions depending on their multimeric association and that thrombin may be critical in their regulation.

## Abstract# 701

Poster Board#/Session: 85-II

**INHIBITION OF ANGIOGENESIS BY PEPTIDES DERIVED FROM KININOGEN.** R.W. Colman, Y. Lin\*, D. Johnson\*, S.A. Mousa\*. Sol Sherry Thrombosis Research Center, Temple University School of Medicine, Philadelphia, PA and DuPont Pharmaceuticals, Wilmington, DE, USA.

Recent studies from our laboratory have shown that cleaved high molecular weight kininogen (HKa) binds specifically, saturably and reversibly and in a zinc-dependent fashion to the human umbilical vein endothelial cell (HUVEC) urokinase receptor (uPAR). The binding of HKa to domain 2/3 of uPAR is inhibited by vitronectin, which binds to the same site, suggesting that HKa is antiadhesive. In addition, HKa is complexed with kallikrein, which cleaves pro-urokinase to urokinase, which confers fibrinolytic activity on HUVEC. We therefore postulated that HKa might be proangiogenic and that peptides from HKa could complete and inhibit angiogenesis by inhibiting the interaction of HKa with uPAR. Migration of HUVEC toward vitronectin and fibronectin was measured in a Boyden chamber fitted with a chemotaxis membrane (8 microns diameter). Adherent cells were permeabilized, stained with rhodamine and the fluorescence quantified. The recombinant polypeptide, HK domain 5 (Lys420-Ser513), at a concentration of 270 nM, inhibited (85%) the migration of HUVEC to vitronectin, but inhibited less than 50% the migration to fibronectin. HUVEC cells were stimulated with basic fibroblast growth factor (b-FGF) for 48 min at 37°C under CO<sub>2</sub> and proliferation measured by the fluorescence enhancement of CyQUANT GR dye bound to nucleic acid of lysed cells. HK domain 5 inhibited HUVEC proliferation (100%) at 270 nM. It also blocked new vessel formation (stimulated by b-FGF) in the chicken chorioallantoic membrane. Deletion mutagenesis indicated that the domain 5 peptide Lys420-Asp474 inhibited HUVEC proliferation 94% at 360 nM. Removal of amino acids Lys420-Asp464 completely abolished the inhibition, indicating that this sequence is critical in the inhibition of HUVEC proliferation. The results suggest that polypeptides from domain 5 of HK may be potent inhibitors of angiogenesis, which could have potential for inhibiting tumor cell metastasis and invasion, diabetic retinopathy (neovascularization) and the abnormal remodeling in atherosclerosis.

## Abstract# 702

Poster Board#/Session: 86-II

**ANGIOSTATIN 4.5: A NATURALLY OCCURRING HUMAN ANGIOGENESIS INHIBITOR.** Gerald A. Soff, Jerome Hong, David Fishman\*, Marlon Kleinman\*, Edward Kaplan\*, David Zagzag\*, Ryan Schultz\*, Deborah Cundiff\*, Susan Park\*, Jan Enghild\*, M. Sharon Stack\*, Stephen Gately\*. Division of Hematology/Oncology, Department of Obstetrics and Gynecology, Northwestern University Medical School, Chicago, IL; Department of Neuropathology, New York University Medical Center, New York, NY; Department of Pathology, Duke University, Durham, NC, USA.

Angiostatin, a proteolytic fragment of plasminogen, inhibits angiogenesis and thereby the growth of primary and metastatic cancers. Although angiostatin has been detected in tumor bearing mice, it has not been previously described in humans. We now report the identification and characterization of an angiostatin isoform in human specimens. Angiostatin4.5 is a 52 kD protein containing kringles

generated in a reaction requiring plasmin catalytic activity in the presence of a free sulfhydryl donor. Western blot assays, antisera raised against the carboxy terminus of angiostatin4.5 as well as a kringle 1-3 antibody detected angiostatin in 100% of human plasma samples from normal (n=10) and cancer patients (n=20), malignant ovarian ascites (n=12), and brain tumor cyst fluids (n=1). The identity of the angiostatin4.5 was further confirmed by its removal by immunoabsorption of plasma with lysine-sepharose. The plasma concentration was approximately 0.1-0.2 µg/ml. Angiostatin4.5, affinity-purified from malignant ascites, was confirmed to be bioactive, inhibiting bFGF-induced vascular endothelial cell proliferation. The previously reported 38-45 kD angiostatin isoform that consists of kringles 1-4 was not detected in any of the specimens. Angiostatin4.5 is the first naturally occurring human angiostatin isoform to be identified and is present in plasma, malignant cyst fluids, and malignant ascites.

## Abstract# 703

Poster Board#/Session: 87-I

**EXPRESSION OF ANGIOPOIETIN 1, ANGIOPOIETIN 2, TIE 1 AND TIE 2 RECEPTORS IN BREAST CANCER.** K. Fujikawa\*, E. Presman\*, C. Higgins\*, Y. Inai\*, L. Varticovski. Department of Biomedical Research, St. Elizabeth Medical Center, Tufts University School of Medicine, Boston, MA, USA.

Several growth factors whose receptors are expressed on endothelial cells have been implicated in regulation of tumor-associated angiogenesis. The progression of many tumors, including breast cancer, is associated with enhanced angiogenesis. Tie1 and Tie2 are endothelial cell receptors of the tyrosine kinase family which regulate angiogenesis in the embryo. Angiopoietin 1 and its naturally-occurring antagonist, Angiopoietin 2, are specific ligands for Tie2 and are required for normal embryonic development of the vascular system. We and others have reported that Tie1 and Tie2 mRNA and proteins are expressed in normal adult endothelial cells and vessels associated with malignant breast lesions. We now examined the expression of Tie1, Tie2, Angiopoietin 1 and Angiopoietin 2 in breast cancer and compared mRNA levels using RT-PCR. Expression of the ligands, Angiopoietin 1 and Angiopoietin 2, and Tie receptors was increased in aggressive breast cancer lesions. In addition, we found that Tie1 and Tie2 proteins are expressed at higher levels in activated microvascular young endothelial cells than in large vessels distal to the malignant lesions. Therefore, Tie1 and Tie2 as the ligands, Angiopoietin 1 and Angiopoietin 2, may participate in progression of breast cancer by enhancing the angiogenic response to the tumor. Detection of Tie receptors and their ligands provides a novel diagnostic tool for identifying tumors with angiogenic potential.

## Abstract# 704

Poster Board#/Session: 88-I

**NEUTRALIZING MONOCLONAL ANTIBODY TO VEGFR-2 (KDR) ABROGATES BASIC FIBROBLAST GROWTH FACTOR INDUCED ANGIOGENESIS.** M. Peichev\*, A.J. Najayer\*, L. Witte\*, D. Hicklin\*, Bohlen\*, R.L. Nachman, M.A.S. Moore, S. Rafii. Hematology Div., Cornell Medical College, ImClone Systems, NY, NY, USA.

Basic Fibroblast growth factor (bFGF) is a cytokine with pleiotropic effects on a variety of cells. Known bFGF functions include stimulation of the migratory chemotaxis, and proliferation of endothelial cells (EC); a process referred to as angiogenesis. To date, the exact mechanism of bFGF function in the regulation of angiogenesis is not well defined. Most *in vitro* studies have used serum-containing media to evaluate the role of bFGF in the regulation of angiogenesis. We have shown that serum contains vascular endothelial growth factor (VEGF) (PNAS 94: 1997). Therefore, the effect of bFGF in these studies may have been mediated through modulation of VEGF receptor-2, (VEGFR-2, KDR), which is the principal specific mitogenic receptor for ECs. Given that expression of KDR is the critical limiting factor in VEGF response, we hypothesized that bFGF may regulate angiogenesis through modulation of KDR expression. Studies analyzing KDR expression have been hampered by the lack of neutralizing anti-KDR MoAb. Therefore, in order to study regulation of KDR, we have developed high affinity as well as neutralizing MoAb to the extracellular domain of KDR. Using FITC conjugated MoAb to KDR we have demonstrated by FACS analysis, that within 1 hour of treatment of human umbilical vein endothelial cells (HUVEC) with bFGF (5 ng/ml), there is a 4.5 fold upregulation of KDR surface expression. In parallel using semi-quantitative RT-PCR we have shown that within 8 hours of incubation of HUVEC with bFGF (5 ng/ml) there is 20 fold increase in the expression of KDR mRNA. Cell cycle analysis using propidium iodide uptake, and cell proliferation as measured by <sup>3</sup>H-thymidine incorporation in HUVECs cultured in serum-free conditions demonstrated that bFGF alone does not support their survival and proliferation. However pretreatment of HUVECs with bFGF (5 ng/ml) followed by VEGF (10 ng/ml), resulted in a 1.5 fold increase in cell number, of which only a small percentage were undergoing apoptosis. In contrast, quantification of early apoptosis by Annexin V expression of the cells that were treated only with bFGF revealed that almost all the cells were in the early stages of apoptosis. Moreover, incubation of HUVECs with neutralizing anti-KDR MoAb (10 µg/ml) blocked the proliferative effect of 10 ng/ml of bFGF by 75 ± 5%. In addition, MoAb to KDR completely inhibited migration of bFGF primed HUVECs through collagen coated Boyden Chambers, as well as endothelial tube formation in matrigel angiogenesis assays. These studies suggest that bFGF induce angiogenesis primarily through upregulation of KDR. Therefore anti-KDR MoAbs may be used as anti-angiogenic factors to treat malignancies that are driven by bFGF and/or VEGF.

VOL 92, NO 10, Supplement 1 (Part 1 of 2) to

NOVEMBER 15, 1998

# BLOOD

*Journal of the American Society of Hematology*

*R. Colman*

American Society of Hematology  
Fortieth Annual Meeting  
December 4-8, 1998  
Miami Beach, FL



# BLOOD

The Journal of  
The American Society of Hematology

*Blood: The Journal of The American Society of Hematology* (ISSN 0006-4971) is published 25 times, in two volumes per year, by W.B. Saunders Company. Dates of issue are the first and the 15th of each month except November in which three issues will be published. W.B. Saunders Company Corporate and Editorial Offices: The Curtis Center, Independence Square West, Philadelphia, PA 19106-3399. Accounting and Circulation Offices: W.B. Saunders Company, Periodicals Dept., 6277 Sea Harbor Dr, Orlando, FL 32887-4800. Periodicals postage paid at Orlando, FL 32862, and at additional mailing offices.

**POSTMASTER:** Send change of address to *Blood: The Journal of The American Society of Hematology*, c/o W.B. Saunders Company, Periodicals Dept., 6277 Sea Harbor Dr, Orlando, FL 32887-4800.

Editorial correspondence should be addressed to:  
Kenneth Kaushansky, MD, Editor-in-Chief

**BLOOD**  
*The Journal of The American Society of Hematology*  
1200 19th Street NW, Suite 300  
Washington, DC 20036-2422

Internet: BLOOD Home Page: [www.bloodjournal.org](http://www.bloodjournal.org). ASH  
Home Page: [www.hematology.org](http://www.hematology.org).

Correspondence regarding subscriptions or change of address should be addressed to W.B. Saunders Company, Periodicals Dept., 6277 Sea Harbor Dr, Orlando, FL 32887-4800.

Customer Service: (800) 654-2452; outside the United States and Canada, (407) 345-4000.

Change of address notices, including both the old and new addresses of the subscriber, should be sent at least one month in advance.

**Yearly subscription rates:** United States and possessions: individual, \$447.00; institution, \$627.00; single issues, \$32.00. All other countries: individual, \$626.00; institution, \$752.00; single issues, \$32.00. For all areas outside the United States and possessions, there is no additional charge for surface delivery. For air mail delivery, add \$144.00. Student and resident: United States and possessions: \$215.00; all other countries: \$588.00. To receive student/resident rate, orders must be accompanied by name of affiliated institution, date of term, and the signature of program/residency coordinator on institution letterhead. Orders will be billed at individual rate until proof of status is received. Back issue and back volume prices are those in current effect. Single issues, both current and back, exist in limited quantities and are offered for sale subject to availability. Back issues sold in conjunction with a subscription are on a prorated basis. Subscriptions are accepted on a calendar year basis. 1997 bound volume price: \$110.00; outside US, please add \$40.00 for postage. To purchase a 1997 bound volume, customer must be a subscriber for 1997. Cumulative Index (1980-1989) price: \$125.00; outside US, please add \$2.25 for surface delivery, or \$8.00 for air mail delivery. Prices are subject to change without notice. Checks should be made payable to W.B. Saunders Company and sent to *Blood: The Journal of The American Society of Hematology*, W.B. Saunders Company, Periodicals Dept., PO Box 628239, Orlando, FL 32862-8239.

Copyright © 1998, The American Society of Hematology. All rights reserved. No part of this publication may be reproduced or transmitted in any form or by any means now or hereafter known, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without permission in writing from the Publisher. Printed in the United States of America.

Correspondance regarding permission to reprint all or part of any article published in this journal should be addressed to Journal Permissions Dept., W.B. Saunders Co., Orlando, FL 32887-4800. Telephone number: (407) 345-2500.

Other correspondence (copyediting, production) should be addressed to:

**BLOOD**  
W.B. Saunders Company  
The Curtis Center  
Independence Square West  
Philadelphia, PA 19106-3399

The appearance of the code at the bottom of the first page of an article in this journal indicates the copyright owner's consent that copies of the article may be made for personal or internal use, or for the personal or internal use of specific clients, for those registered with the Copyright Clearance Center, Inc. (222 Rosewood Drive, Danvers, MA 01923; (978) 750-8400; [www.copyright.com](http://www.copyright.com)). This consent is given on the condition that the copier pay the stated per-copy fee for that article through the Copyright Clearance Center, Inc. for copying beyond that permitted by Sections 107 or 108 of the US Copyright Law. This consent does not extend to other kinds of copying, such as copying for general distribution, for advertising or promotional purposes, for creating new collective works, or for resale. Absence of the code indicates that the material may not be processed through the Copyright Clearance Center, Inc.

**Advertising Representative:** Cunningham Associates, 180 Old Tappan Rd, Old Tappan, NJ 07675, telephone (201) 767-4170, fax (201) 767-8065.

The ideas and opinions expressed in *Blood* do not necessarily reflect those of The American Society of Hematology, the Editor or the Publisher. Publication of an advertisement or other product mention in *Blood* should not be construed as an endorsement of the product or the manufacturer's claims. Readers are encouraged to contact the manufacturer with any questions about the features or limitations of the products mentioned. Neither The American Society of Hematology nor the Publisher assumes any responsibility for any injury and/or damage to persons or property arising out of or related to any use of the material contained in this periodical. The reader is advised to check the appropriate medical literature and the product information currently provided by the manufacturer of each product to be administered to verify the dosage, the method and duration of administration, or contraindications. It is the responsibility of the treating physician or other health care professional, relying on independent experience and knowledge of the patient, to determine drug dosages and the best treatment for the patient.



PROGRAM OF THE 40th ANNUAL MEETING OF  
**THE AMERICAN SOCIETY OF HEMATOLOGY**

---

*December 4-8, 1998*

*Miami Beach, Florida*

---

---

Abstract management services by PRISM Productions, Inc., Westerville, Ohio  
Printed in the USA  
COPYRIGHT© 1998, THE AMERICAN SOCIETY OF HEMATOLOGY, WASHINGTON, DC